

Fluctuations in Relativistic Causal Hydrodynamics

Avdhesh Kumar, Jitesh R. Bhatt and Ananta P. Mishra

Physical Research Laboratory, Navrangpura, Ahmedabad 380 009, India

E-mail addresses: avdhesh@prl.res.in, jeet@prl.res.in, apmishra@prl.res.in

ABSTRACT: Formalism to calculate the hydrodynamic fluctuations by applying the Onsager theory to the relativistic Navier-Stokes equation is already known. In this work, we calculate hydrodynamic-fluctuations in the framework of the causal hydrodynamics of Müller, Israel and Stewart and the other related approaches. We show that expressions for the Onsager-coefficients and the correlation-functions have forms similar to the ones obtained by using the relativistic Navier-Stokes equation. However, spatio-temporal evolution of the correlation functions obtained using MIS and the other causal theories can be significantly different from the correlation functions obtained using the Navier-Stokes equation. Finally, as an illustrative example, we numerically evaluate the correlation-functions using the one dimensional expanding boost-invariant (Bjorken) flows and compare the correlation-functions obtained using the various hydrodynamic approaches.

Contents

1. Introduction	1
2. Fluctuations and correlations in Hydrodynamics	2
2.1 Equations for dissipative fluxes in Landau-Lifshitz frame and fluctuation correlations in MIS	4
2.2 Equations for dissipative fluxes in Eckart frame and fluctuation correlations in MIS	6
2.3 Equations for dissipative fluxes in Landau-Lifshitz frame and fluctuation correlations for other Hydrodynamic models	8
2.3.1 Third order hydrodynamics	8
2.3.2 JBP hydrodynamics	9
2.3.3 DKR hydrodynamics	10
2.3.4 Conformal viscous hydrodynamics	12
3. Calculation of correlation functions in boost-invariant Hydrodynamics	12
4. Results and Discussions	14
5. Conclusions	18
6. Acknowledgements	18

1. Introduction

A study of fluctuations in continuous media is of great interest in physics and it can provide a link between the macroscopic and microscopic points of view. A macroscopic theory such as hydrodynamics provides a simplest possible description of a complicated many-body system in terms of space-time evolution of the mean or averaged quantities like energy density, pressure, averaged flow velocity etc. The fluctuation theory studies small deviations from the mean behavior and it can help in calculating correlation functions for the macroscopic variables [1, 2]. In context of relativistic hydrodynamics, results of the fluctuation-dissipation theorem has been studied in Refs. [3, 4]. In Ref. [3] authors have studied the fluctuation in the contexts of general-relativistic Navier-Stokes theory. A more general framework of hydrodynamics described as the divergence type theory(DTT) [5] was considered in Ref. [4]. It ought to be noted that recently in an interesting work in Ref. [6], the authors have applied the results of fluctuation-dissipation theorem to the relativistic *Navier-Stokes* theory of hydrodynamics and calculated the two particles correlators for

the one-dimensional hydrodynamics (Bjorken) flow relevant for the relativistic heavy-ion collision experiments at RHIC and LHC. The authors obtained several analytical results for two particle correlation functions. Further, in Ref. [7], the authors have studied the effect of thermal conductivity on the correlation function using the Bjorken-flow. It should be noted here that it is well-known that relativistic Navier-Stokes theory exhibit acausal behavior and it can give rise to unphysical instabilities [8]. However the causality can be restored if the terms with higher orders are included in the hydrodynamics as indicated by the Maxwell-Cattaneo law [9]. Indeed these issues do not arise in the second-order causal hydrodynamics theory developed by Müller, Israel and Stewart (MIS) [10]. The Navier-Stokes equations can be derived from the second law of thermodynamics $\partial_\mu S^\mu \geq 0$, where S^μ denotes the equilibrium entropy current [2]. However, it is not necessary for an out-of-equilibrium fluid to have an equilibrium entropy current [9]. In MIS hydrodynamics out-of-equilibrium current can have contributions from dissipative processes like the effect of viscosity and the heat conduction. This has an interesting analogy with the irreversible thermodynamics [11, 12]. Further the MIS hydrodynamics has been extensively applied to study the relativistic heavy-ion collisions [9, 13–15] and also in cosmology [16]. Later this formalism was extended to include the effect of third order terms in the gradient expansion [17]. Recently, it has been shown that the derivation of the MIS equations from the underlying kinetic equation may not be unique, there may exist a more general set of hydrodynamic equations which may allow one to obtain MIS equations as a special case [18, 19].

Finally, it should be mentioned here that although the divergent type theory (DTT) of relativistic fluid of Geroch-Lindblom [5] allows for a consistent proof of causality and stability of its solutions, it is far from direct thermodynamic intuition. Moreover, the connection between the DTTs and MIS or other causal hydrodynamics theories is not yet clearly established.

In this work we apply the fluctuation-dissipation theorem to MIS equations and also to the hydrodynamics models developed by Denicol, Koide and Rischke (DKR) [19], Jaiswal, Bhalerao and Pal (JBP) [18] and other models based on MIS approach [10, 17, 20]. Further, we apply the results obtained to study the hydrodynamical evolution using 1+0 dimensional Bjorken flow. In particular, we calculate the correlators using the Onsager coefficients for various relativistic hydrodynamical theories. We believe that such study may be important in choosing the appropriate version of the causal hydrodynamics from the variety of the options available.

2. Fluctuations and correlations in Hydrodynamics

In thermodynamic equilibrium entropy of the system S which is function of the additive quantities x_k becomes maximum. In equilibrium, S satisfies the condition $X_k = -\frac{\partial S}{\partial x_k} = 0$. However, when the system is slightly away from the equilibrium the generalized forces $X_k \neq 0$ and $\frac{dx_i}{dt} = -\gamma_{ik}X_k + \xi_i$, the summation convention is implied, describes the flux associated with the quantity x_i , here ξ_i are the random forces or the noise term and γ_{ik} are the Onsager coefficients. The Onsager reciprocity relations imply that $\gamma_{ik} = \gamma_{ki}$. In this

phenomenological theory time rate of change of the total entropy, $\frac{dS}{dt}$ can be given by,

$$\frac{dS}{dt} = -\frac{dx_i}{dt}X_i, \quad (2.1)$$

which can also be written as,

$$\frac{dS}{dt} = \gamma_{ik}X_kX_i - \xi_iX_i. \quad (2.2)$$

Noises are correlated and the correlation is given by the formula,

$$\langle \xi_i(t_1)\xi_k(t_2) \rangle = (\gamma_{ik} + \gamma_{ki})\delta(t_1 - t_2). \quad (2.3)$$

The correlation functions can be found once γ_{ij} are known [2, 6]. In order to find out γ_{ij} , one needs to know the $\frac{dS}{dt}$ for the hydrodynamical theory and identify the generalized forces and fluxes. The expression for entropy rate $\frac{dS}{dt}$ can be found either by using equations of hydrodynamics together with the thermodynamic relations or from kinetic theory [10]. In our work we are using the former approach.

We start with the expressions for the energy momentum tensor $T^{\mu\nu}$ and the current-density J_B^μ for a viscous fluid,

$$T^{\mu\nu} = T_{id}^{\mu\nu} + \Delta T^{\mu\nu} + S^{\mu\nu}, \quad (2.4)$$

$$J_B^\mu = n_B u^\mu + \nu^\mu + I^\mu, \quad (2.5)$$

where, $T_{id}^{\mu\nu} = \epsilon u^\mu u^\nu - p \Delta^{\mu\nu}$ is the ideal part of the energy momentum tensor with $\Delta^{\mu\nu} = g^{\mu\nu} - u^\mu u^\nu$, and ϵ , p , u^μ are the local energy density, pressure and fluid flow four-velocity respectively. It is to be noted that $u^\mu u_\mu = 1$ and $g_{\mu\nu} = \text{diag}(+, -, -, -)$. $\Delta T^{\mu\nu} = \Delta T_{vis}^{\mu\nu} + \Delta T_{heat}^{\mu\nu}$ with $\Delta T_{vis}^{\mu\nu} = \pi^{\mu\nu} - \Delta^{\mu\nu} \Pi$ and $\Delta T_{heat}^{\mu\nu} = W^\mu u^\nu + W^\nu u^\mu$, is the dissipative part of the energy momentum tensor and $S^{\mu\nu}$ is the stochastic term arising due to the local thermal fluctuations [6]. Similarly, ν^μ and I^μ denote the dissipative(non-equilibrium) and the stochastic terms in baryon current density respectively. n_B denotes the net density of baryon number in local rest frame. $W^\mu = q^\mu + h\nu^\mu$ is the energy flow in local rest frame, $h = (\epsilon + p)/n$ is the enthalpy per particle and $\nu^\mu = \Delta^{\mu\nu} J_B^\nu$ is the baryon number flow in the local rest frame. For the dissipative fluxes one can always require the following relations to hold $u_\mu \pi^{\mu\nu} = 0$, $\pi_\alpha^\alpha = 0$, $\Delta_{\mu\nu} \pi^{\mu\nu} = 0$, $u_\mu W^\mu = 0$, $u_\mu \nu^\mu = 0$, $u_\mu q^\mu = 0$.

The relevant conservation equations for the hydrodynamics can be written as,

$$\partial_\mu J_B^\mu = Dn_B + n_B \nabla_\mu u^\mu + \partial_\mu \nu^\mu = 0, \quad (2.6)$$

$$u_\nu \partial_\mu T^{\mu\nu} = D\epsilon + (\epsilon + p + \Pi) \nabla_\mu u^\mu - \pi_{\mu\nu} \nabla^{(\mu} u^{\nu)} + \nabla_\mu W^\mu - 2W^\mu Du_\mu = 0, \quad (2.7)$$

$$\begin{aligned} \Delta_\nu^\alpha \partial_\mu T^{\mu\nu} &= (\epsilon + p + \Pi) Du^\alpha - \nabla^\alpha (p + \Pi) + \Delta^{\alpha\nu} \nabla^\sigma \pi_{\nu\sigma} - \pi^{\alpha\nu} Du_\nu \\ &\quad + \Delta^{\alpha\nu} DW_\nu + 2W^{(\alpha} \nabla_\nu u^{\nu)} = 0, \end{aligned} \quad (2.8)$$

where, $D = u^\mu \partial_\mu$ and $\nabla^\mu = \Delta^{\mu\nu} \partial_\nu$. There are only 5 equations written above and 14 unknowns n_B , ϵ , Π , W^μ , $\pi^{\mu\nu}$ and u^μ . Therefore, 9 additional equations for dissipative fluxes are required to obtain the hydrodynamical solution.

There are two popular choices for u^μ : In Landau-Lifshitz frame u^μ is parallel to the energy-flow and $W^\mu = 0$ which implies that $q^\mu = -h\nu^\mu$. Another choice is the Eckart-frame, where u^μ can be parallel to J_B^μ and $\nu^\mu = 0$ and this would imply $W^\mu = q^\mu$. Now we shall obtain the 9 additional equations and fluctuation correlations in Landau-Lifshitz and Eckart frame respectively.

2.1 Equations for dissipative fluxes in Landau-Lifshitz frame and fluctuation correlations in MIS

In order to derive the 9 additional equations one needs the expression for the out-of-equilibrium entropy four-current. In Landau-Lifshitz frame the expression for the non-equilibrium entropy four-current is given in literature [10, 14, 15] and is as follows,

$$S^\mu = su^\mu - \frac{\mu_B}{T}\nu^\mu - \left(\beta_0\Pi^2 - \beta_1q_\nu q^\nu + \beta_2\pi_{\nu\lambda}\pi^{\nu\lambda}\right)\frac{u^\mu}{2T} - \frac{\alpha_0\Pi q^\mu}{T} + \frac{\alpha_1\pi^{\mu\nu}q_\nu}{T}. \quad (2.9)$$

The divergence of the non-equilibrium entropy four-current (Eq.(2.9)) using Eqs.(2.6-2.7) and the thermodynamic relations $d\epsilon = Tds - \mu dn$ and $\epsilon + p = Ts - \mu n$ can be written as follows,

$$\begin{aligned} T\partial_\mu S^\mu = & -\Pi \left[\partial_\mu u^\mu + \beta_0\dot{\Pi} + \frac{1}{2}T\partial_\mu \left(\frac{\beta_0}{T}u^\mu \right) \Pi + \alpha_0\nabla_\mu q^\mu \right] \\ & - q^\mu \left[-h^{-1}T\nabla_\mu \left(\frac{\mu}{T} \right) - \beta_1\dot{q}_\mu - \frac{1}{2}T\partial_\nu \left(\frac{\beta_1}{T}u^\nu \right) q_\mu - \alpha_1\partial_\nu \pi_\mu^\nu + \alpha_0\partial_\mu \Pi \right] \\ & + \pi^{\mu\nu} \left[\sigma_{\mu\nu} - \beta_2\dot{\pi}_{\mu\nu} - \frac{1}{2}T\partial_\lambda \left(\frac{\beta_2}{T}u^\lambda \right) \pi_{\mu\nu} + \alpha_1\nabla_{\langle\nu} q_{\mu\rangle} \right], \end{aligned} \quad (2.10)$$

where we have used the notation $\dot{F} = DF$. According to the second law of thermodynamics we must have $T\partial_\mu S^\mu \geq 0$. This inequality will be satisfied if Π, q_μ and $\pi^{\mu\nu}$ satisfy the following equations,

$$\left[\partial_\mu u^\mu + \beta_0\dot{\Pi} + \frac{1}{2}T\partial_\mu \left(\frac{\beta_0}{T}u^\mu \right) \Pi + \alpha_0\nabla_\mu q^\mu \right] = -\frac{\Pi}{\zeta}, \quad (2.11)$$

$$\left[-h^{-1}T\nabla_\mu \left(\frac{\mu}{T} \right) - \beta_1\dot{q}_\mu - \frac{1}{2}T\partial_\nu \left(\frac{\beta_1}{T}u^\nu \right) q_\mu - \alpha_1\partial_\nu \pi_\mu^\nu + \alpha_0\partial_\mu \Pi \right] = \frac{q_\mu}{\lambda T}, \quad (2.12)$$

$$\left[\sigma_{\mu\nu} - \beta_2\dot{\pi}_{\mu\nu} - \frac{1}{2}T\partial_\lambda \left(\frac{\beta_2}{T}u^\lambda \right) \pi_{\mu\nu} + \alpha_1\nabla_{\langle\nu} q_{\mu\rangle} \right] = \frac{\pi_{\mu\nu}}{2\eta}. \quad (2.13)$$

Here we note that the term with factor 1/2 on the left hand side of equations (2.11-2.13) are generally ignored by arguing that the gradient of thermodynamic quantities are small [15, 21]. Thus, Eqs.(2.11-2.13) can be written as,

$$\tau_\Pi\dot{\Pi} + \Pi = -\zeta\nabla_\alpha u^\alpha - l_{\Pi q}\nabla_\mu q^\mu, \quad (2.14)$$

$$\tau_q\dot{q}_\mu + q_\mu = -\lambda T^2 h^{-1}\nabla_\mu \left(\frac{\mu}{T} \right) + l_{q\Pi}\nabla_\mu \Pi - l_{q\pi}\nabla_\nu \pi_\mu^\nu, \quad (2.15)$$

$$\tau_\pi\dot{\pi}_{\mu\nu} + \pi_{\mu\nu} = 2\eta\sigma_{\mu\nu} + l_{\pi q}\nabla_{\langle\mu} q_{\nu\rangle}. \quad (2.16)$$

Henceforth, we call Eqs.(2.14-2.16) as MIS equations. Here, $\tau_\Pi = \zeta\beta_0$, $\tau_q = \lambda T\beta_1$, $\tau_\pi = 2\eta\beta_2$ are identified as the relaxation times and $l_{\Pi q} = \zeta\alpha_0$, $l_{q\Pi} = \lambda T\alpha_0$, $l_{q\pi} = \lambda T\alpha_1$, $l_{\pi q} = 2\eta\alpha_1$ as coupling constants. These 9 equations for the dissipative fluxes combined with the Eqs.(2.6-2.8) form complete set of the hydrodynamic equations. Note that the limit $\tau_\Pi, \tau_q, \tau_\pi, l_{\Pi q}, l_{q\Pi}, l_{q\pi}, l_{\pi q} \rightarrow 0$ is the first order limit which correspond to the Navier-Stokes case.

Eq.(2.10) can be written as,

$$T\partial_\mu S^\mu = \frac{\Pi^2}{\zeta} - \frac{q^\mu q_\mu}{\lambda T} + \frac{\pi^{\mu\nu}\pi_{\mu\nu}}{2\eta} \geq 0. \quad (2.17)$$

Here, $q^\mu q_\mu < 0$ [13]. Now using the identity $\Delta^{\mu\nu}\Delta_{\mu\nu} = 3$ and the condition $\Delta_{\mu\nu}\pi^{\mu\nu} = 0$, one can write Eq.(2.17) as,

$$\partial_\mu S^\mu = \frac{\Delta T_{vis}^{\mu\nu}}{T} \left(\frac{\pi_{\mu\nu}}{2\eta} - \frac{\Delta_{\mu\nu}\Pi}{3\zeta} \right) - \frac{q^\mu q_\mu}{\lambda T^2}. \quad (2.18)$$

Upon integrating over the whole volume Eq.(2.18) can be written as,

$$\frac{dS}{dt} = \int d^3x \left[\frac{\Delta T_{vis}^{\mu\nu}}{T} \left(\frac{\pi_{\mu\nu}}{2\eta} - \frac{\Delta_{\mu\nu}\Pi}{3\zeta} \right) - \frac{q^\mu q_\mu}{\lambda T^2} \right]. \quad (2.19)$$

Following identification between the phenomenological variables $(\dot{x}_1, \dot{x}_2)$ and the hydrodynamical variables can be made [6],

$$\dot{x}_1 \rightarrow \Delta T_{vis}^{\mu\nu}, \quad \dot{x}_2 \rightarrow q^\mu. \quad (2.20)$$

A comparison of Eq.(2.19) with the phenomenological equation Eq.(2.1) will give,

$$\begin{aligned} X_1 &= -\frac{1}{T} \left(\frac{\pi_{\mu\nu}}{2\eta} - \frac{\Delta_{\mu\nu}\Pi}{3\zeta} \right) \Delta V, \\ X_2 &= \frac{q_\mu}{\lambda T^2} \Delta V. \end{aligned} \quad (2.21)$$

Now neglecting the stochastic term in Eq.(2.2) and comparing it with Eq.(2.19) one can get,

$$X_1 = -\Delta T_{vis}^{\mu\nu}, \quad (2.22)$$

$$\gamma_{22} X_2 = -q^\mu, \quad (2.23)$$

$$\gamma_{12} = \gamma_{21} = 0 \quad (2.24)$$

The coefficients γ_{12} and γ_{21} are zero, because dissipative fluxes due to heat and viscosity are treated to be independent. Coefficients γ_{11} and γ_{22} are the rank-four tensor and they can be parameterized as follows,

$$\gamma_{11} = \left[A\Delta^{\mu\nu\alpha\beta} + B\Delta^{\mu\nu}\Delta^{\alpha\beta} \right] \frac{1}{\Delta V}, \quad \gamma_{22} = \frac{C\Delta^{\mu\nu}}{\Delta V}, \quad (2.25)$$

where, $\Delta^{\mu\nu\alpha\beta} = \Delta^{\mu\alpha}\Delta^{\nu\beta} - \frac{1}{3}\Delta^{\mu\nu}\Delta^{\alpha\beta}$. Now using Eqs.(2.22, 2.23) one can determine the coefficients $A = 2\eta T$, $B = \zeta T$ and $C = -\lambda T^2$. Thus one can write,

$$\gamma_{11} = 2T \left[\left(\eta \Delta^{\mu\alpha}\Delta^{\nu\beta} - \frac{1}{3}\eta \Delta^{\mu\nu}\Delta^{\alpha\beta} \right) + \frac{1}{2}\zeta \Delta^{\mu\nu}\Delta^{\alpha\beta} \right] \frac{1}{\Delta V}, \quad \gamma_{22} = -\frac{\lambda T^2 \Delta^{\mu\nu}}{\Delta V}. \quad (2.26)$$

From above expression of γ_{11} one can see that there is a additive contribution of shear and bulk viscosity i.e one can write it as $\gamma_{11} = (\gamma_{11})_\eta + (\gamma_{11})_\zeta$.

Now following Eq.(2.2), the correlation functions can be written as,

$$\langle S_{vis}^{\mu\nu}(x_1) S_{vis}^{\alpha\beta}(x_2) \rangle = 2T \left[\eta (\Delta^{\mu\alpha}\Delta^{\nu\beta} + \Delta^{\mu\beta}\Delta^{\nu\alpha}) + (\zeta - \frac{2}{3}\eta) \Delta^{\mu\nu}\Delta^{\alpha\beta} \right] \delta(x_1 - x_2), \quad (2.27)$$

$$\langle I^\mu(x_1) I^\nu(x_2) \rangle = -2\lambda T^2 \Delta^{\mu\nu} \delta(x_1 - x_2), \quad (2.28)$$

$$\langle S_{vis}^{\mu\nu}(x_1) I^\alpha(x_2) \rangle = 0. \quad (2.29)$$

These are the stochastic or noise auto-correlation functions for the MIS hydrodynamics in the Landau-Lifshitz frame.

2.2 Equations for dissipative fluxes in Eckart frame and fluctuation correlations in MIS

In Eckart frame the expression for the entropy four-current [10, 13, 14] can be written as,

$$S^\mu = s u^\mu + \frac{q_\mu}{T} - \left(\beta_0 \Pi^2 - \bar{\beta}_1 q_\nu q^\nu + \beta_2 \pi_{\nu\lambda} \pi^{\nu\lambda} \right) \frac{u^\mu}{2T} - \frac{\bar{\alpha}_0 \Pi q^\mu}{T} + \frac{\bar{\alpha}_1 \pi^{\mu\nu} q_\nu}{T}. \quad (2.30)$$

Note that here coefficients β_0, β_2 are same as in Landau-Lifshitz case while the coefficients $\bar{\alpha}_i$ and $\bar{\beta}_1$ are given as, $\bar{\alpha}_i = \alpha_i + \frac{1}{\epsilon+p}$ and $\bar{\beta}_1 = \beta_1 + \frac{1}{\epsilon+p}$, where α_i, β_1 are the coefficients in Landau-Lifshitz frame. Next, the divergence of the non-equilibrium entropy four-current (Eq.(2.30)) using Eqs.(2.6-2.7) and the thermodynamic relations $d\epsilon = Tds - \mu dn$ and $\epsilon + p = Ts - \mu n$ can be written as follows,

$$\begin{aligned} T\partial_\mu S^\mu = & -\Pi \left[\partial_\mu u^\mu + \beta_0 \dot{\Pi} + \frac{1}{2}T\partial_\mu \left(\frac{\beta_0}{T} u^\mu \right) \Pi + \bar{\alpha}_0 \nabla_\mu q^\mu \right] \\ & - q^\mu \left[\nabla_\mu \ln T - \dot{u}_\mu - \bar{\beta}_1 \dot{q}_\mu - \frac{1}{2}T\partial_\nu \left(\frac{\bar{\beta}_1}{T} u^\nu \right) q_\mu - \bar{\alpha}_1 \partial_\nu \pi_\mu^\nu + \bar{\alpha}_0 \partial_\mu \Pi \right] \\ & + \pi^{\mu\nu} \left[\sigma_{\mu\nu} - \beta_2 \dot{\pi}_{\mu\nu} - \frac{1}{2}T\partial_\lambda \left(\frac{\beta_2}{T} u^\lambda \right) \pi_{\mu\nu} + \bar{\alpha}_1 \nabla_{\langle \nu} q_{\mu \rangle} \right]. \end{aligned} \quad (2.31)$$

In order to have $T\partial_\mu S^\mu \geq 0$ we must have,

$$\left[\partial_\mu u^\mu + \beta_0 \dot{\Pi} + \frac{1}{2}T\partial_\mu \left(\frac{\beta_0}{T} u^\mu \right) \Pi + \bar{\alpha}_0 \nabla_\mu q^\mu \right] = -\frac{\Pi}{\zeta}, \quad (2.32)$$

$$\left[\nabla_\mu \ln T - \dot{u}_\mu - \bar{\beta}_1 \dot{q}_\mu - \frac{1}{2}T\partial_\nu \left(\frac{\bar{\beta}_1}{T} u^\nu \right) q_\mu - \bar{\alpha}_1 \partial_\nu \pi_\mu^\nu + \bar{\alpha}_0 \partial_\mu \Pi \right] = \frac{q_\mu}{\lambda T}, \quad (2.33)$$

$$\left[\sigma_{\mu\nu} - \beta_2 \dot{\pi}_{\mu\nu} - \frac{1}{2}T\partial_\lambda \left(\frac{\beta_2}{T} u^\lambda \right) \pi_{\mu\nu} + \bar{\alpha}_1 \nabla_{\langle \nu} q_{\mu \rangle} \right] = \frac{\pi_{\mu\nu}}{2\eta}. \quad (2.34)$$

Thus one can write Eq.(2.31) as,

$$T\partial_\mu S^\mu = \frac{\Pi^2}{\zeta} - \frac{q^\mu q_\mu}{\lambda T} + \frac{\pi^{\mu\nu}\pi_{\mu\nu}}{2\eta} \geq 0, \quad (2.35)$$

which can also be written as,

$$T\partial_\mu S^\mu = \Delta T^{\mu\nu} \left[\frac{\pi_{\mu\nu}}{2\eta} - \frac{\Delta_{\mu\nu}\Pi}{3\zeta} - \frac{1}{2\lambda T}(u_\nu q_\mu + u_\mu q_\nu) \right]. \quad (2.36)$$

Upon integrating over whole volume Eq.(2.36) can be written as,

$$\frac{dS}{dt} = \int d^3x \frac{\Delta T^{\mu\nu}}{T} \left[\frac{\pi_{\mu\nu}}{2\eta} - \frac{\Delta_{\mu\nu}\Pi}{3\zeta} - \frac{1}{2\lambda T}(u_\nu q_\mu + u_\mu q_\nu) \right], \quad (2.37)$$

which can be rearranged in the following form,

$$\frac{dS}{dt} = \int d^3x \left[\frac{\Delta T_{vis}^{\mu\nu}}{T} \left(\frac{\pi_{\mu\nu}}{2\eta} - \frac{\Delta_{\mu\nu}\Pi}{3\zeta} \right) + \frac{\Delta T_{heat}^{\mu\nu}}{T} \left[\left(\frac{\pi_{\mu\nu}}{2\eta} - \frac{\Delta_{\mu\nu}\Pi}{3\zeta} \right) - \frac{1}{2\lambda T}(u_\nu q_\mu + u_\mu q_\nu) \right] \right]. \quad (2.38)$$

In this case also one can make the identifications as before,

$$\dot{x}_1 \rightarrow \Delta T_{vis}^{\mu\nu}, \quad \dot{x}_2 \rightarrow \Delta T_{heat}^{\mu\nu}. \quad (2.39)$$

The comparison between Eqs. (2.38) and (2.1) will give,

$$\begin{aligned} X_1 &= -\frac{1}{T} \left(\frac{\pi_{\mu\nu}}{2\eta} - \frac{\Delta_{\mu\nu}\Pi}{3\zeta} \right) \Delta V, \\ X_2 &= -\frac{1}{T} \left[\left(\frac{\pi_{\mu\nu}}{2\eta} - \frac{\Delta_{\mu\nu}\Pi}{3\zeta} \right) - \frac{1}{2\lambda T}(u_\nu q_\mu + u_\mu q_\nu) \right] \Delta V. \end{aligned} \quad (2.40)$$

Again neglecting the stochastic term in Eq.(2.2) and comparing it with Eq.(2.38) one can get,

$$\gamma_{11} X_1 = -\Delta T_{vis}^{\mu\nu}, \quad (2.41)$$

$$\gamma_{22} X_2 = -\Delta T_{heat}^{\mu\nu}, \quad (2.42)$$

$$\gamma_{12} = \gamma_{21} = 0. \quad (2.43)$$

Now parameterizing γ_{11} and γ_{22} as follows,

$$\gamma_{11} = \left[A\Delta^{\mu\nu\alpha\beta} + B\Delta^{\mu\nu}\Delta^{\alpha\beta} \right] \frac{1}{\Delta V}, \quad \gamma_{22} = \left[\bar{A}\Delta^{\mu\alpha}u^\nu u^\beta + \bar{B}\Delta^{\nu\beta}u^\mu u^\alpha \right] \frac{1}{\Delta V}, \quad (2.44)$$

since we know the form of (X_1, X_2) and $(\Delta T_{vis}^{\mu\nu}, \Delta T_{heat}^{\mu\nu})$ therefore using the above parameterizations of γ_{11} and γ_{22} and Eqs.(2.41-2.42), one can determine the coefficients $A = 2\eta T$, $B = \zeta T$ and $\bar{A} = \bar{B} = -2\lambda T^2$. Therefore the Onsager coefficients are,

$$\gamma_{11} = 2T \left[\left(\eta\Delta^{\mu\alpha}\Delta^{\nu\beta} - \frac{1}{3}\eta\Delta^{\mu\nu}\Delta^{\alpha\beta} \right) + \frac{1}{2}\zeta\Delta^{\mu\nu}\Delta^{\alpha\beta} \right] \frac{1}{\Delta V}, \quad (2.45)$$

$$\gamma_{22} = -2\lambda T^2 \left[\Delta^{\mu\alpha}u^\nu u^\beta + \Delta^{\nu\beta}u^\mu u^\alpha \right] \frac{1}{\Delta V}. \quad (2.46)$$

Thus one can write the correlation functions using Eq.(2.3) as,

$$\langle S_{vis}^{\mu\nu}(x_1) S_{vis}^{\alpha\beta}(x_2) \rangle = 2T \left[\eta(\Delta^{\mu\alpha} \Delta^{\nu\beta} + \Delta^{\mu\beta} \Delta^{\nu\alpha}) + (\zeta - \frac{2}{3}\eta) \Delta^{\mu\nu} \Delta^{\alpha\beta} \right] \delta(x_1 - x_2), \quad (2.47)$$

$$\begin{aligned} \langle S_{heat}^{\mu\nu}(x_1) S_{heat}^{\alpha\beta}(x_2) \rangle = & -2\lambda T^2 \left[\Delta^{\mu\alpha} u^\nu u^\beta + \Delta^{\nu\beta} u^\mu u^\alpha + \Delta^{\mu\beta} u^\nu u^\alpha \right. \\ & \left. + \Delta^{\nu\alpha} u^\mu u^\beta \right] \delta(x_1 - x_2), \end{aligned} \quad (2.48)$$

$$\langle S_{vis}^{\mu\nu}(x_1) S_{heat}^{\alpha\beta}(x_2) \rangle = 0. \quad (2.49)$$

The form of these correlations is very similar to the correlations obtained for the relativistic Navier-Stokes case [6]. However, the evolution of the correlations can be very different as demonstrated later.

2.3 Equations for dissipative fluxes in Landau-Lifshitz frame and fluctuation correlations for other Hydrodynamic models

In this section we consider some of the interesting alternate approaches to the causal MIS hydrodynamics and some of its extensions.

2.3.1 Third order hydrodynamics

In the Ref. [17] third order corrections to the MIS hydrodynamics was considered when the effect of bulk-viscosity and heat-flux were absent. In this case expression for the non-equilibrium entropy four-current can be written as,

$$S^\mu = s u^\mu - \frac{\beta_2}{2T} \pi_{\alpha\beta} \Pi^{\alpha\beta} u^\mu + \alpha \frac{\beta_2^2}{T} \pi_{\alpha\beta} \pi_\sigma^\alpha \pi^{\beta\sigma} u^\mu, \quad (2.50)$$

where, α is a new dimensionless coefficient and it is assumed to be a constant. The last term on the right hand side of the above equation represents the third order correction to the equation of entropy. In order to fulfill the requirement of maximal entropy at equilibrium, the third order term must satisfy the condition $\alpha \frac{\beta_2^2}{T} \pi_{\alpha\beta} \pi_\sigma^\alpha \pi^{\beta\sigma} u^\mu \leq 0$. Divergence of the entropy four-current can be written as,

$$\begin{aligned} \partial_\mu S^\mu = & \frac{1}{T} \pi_{\alpha\beta} \sigma^{\alpha\beta} - \pi_{\alpha\beta} \pi^{\alpha\beta} \partial_\mu \left(\frac{\beta_2}{2T} u^\mu \right) - \frac{\beta_2}{T} \pi_{\alpha\beta} \dot{\pi}^{\alpha\beta} \\ & + \alpha \partial_\mu \left(\frac{\beta_2^2}{T} u^\mu \right) \pi_{\alpha\beta} \pi_\sigma^\alpha \pi^{\beta\sigma} + 3\tau_\pi \theta \alpha \frac{\beta_2^2}{T} \pi_{\alpha\beta} \pi_\sigma^\alpha \dot{\pi}^{\beta\sigma} \geq 0. \end{aligned} \quad (2.51)$$

Here, the Knudsen number($=\tau_\pi \theta$) is required to satisfy the condition $\tau_\pi \theta \ll 1$ for the validity of hydrodynamic approach. For the condition $T \partial_\mu S^\mu \geq 0$ to be satisfied one must have,

$$\partial_\mu s^\mu = \frac{1}{2\eta T} \pi^{\mu\nu} \pi_{\mu\nu}, \quad (2.52)$$

which implies that the form of shear viscous tensor $\pi^{\alpha\beta}$ should be given by

$$\pi^{\alpha\beta} = 2\eta T \left[\frac{1}{T} \sigma^{\alpha\beta} - \pi^{\alpha\beta} \partial_\mu \left(\frac{\beta_2}{2T} u^\mu \right) - \frac{\beta_2}{T} \dot{\pi}^{\alpha\beta} + \alpha \partial_\mu \left(\frac{\beta_2^2}{T} u^\mu \right) \pi_\sigma^\alpha \pi^{\beta\sigma} + 3\tau_\pi \theta \alpha \frac{\beta_2^2}{T} \pi_\sigma^\alpha \dot{\pi}^{\beta\sigma} \right]. \quad (2.53)$$

Now starting from Eq.(2.52) and following similar prescription to determine the Onsager coefficient as done in second-order MIS hydrodynamics one can write

$$\dot{x} = \pi^{\mu\nu}, \quad X = -\frac{1}{2\eta T} \pi^{\mu\nu} \Delta V, \quad (2.54)$$

and the Onsager coefficient,

$$\gamma = 2\eta T \left[\Delta^{\mu\alpha} \Delta^{\nu\beta} - \frac{1}{3} \Delta^{\mu\nu} \Delta^{\alpha\beta} \right] \frac{1}{\Delta V}. \quad (2.55)$$

The viscous correlation function can be written as,

$$\langle S_{vis}^{\mu\nu}(x_1) S_{vis}^{\alpha\beta}(x_2) \rangle = 2T \left[\eta (\Delta^{\mu\alpha} \Delta^{\nu\beta} + \Delta^{\mu\beta} \Delta^{\nu\alpha}) - \frac{2}{3} \eta \Delta^{\mu\nu} \Delta^{\alpha\beta} \right] \delta(x_1 - x_2). \quad (2.56)$$

One can notice that this expression is same as the one obtained using the second-order theory with $\Pi = 0$.

2.3.2 JBP hydrodynamics

In the Ref [18] the authors have constructed the expression for the entropy four-current S^μ generalized from the Boltzmann's H-function and find out the expression for its divergence as,

$$\begin{aligned} \partial_\mu S^\mu = & -\frac{\Pi}{T} \left[\theta + \beta_0 \dot{\Pi} + \beta_{\Pi\Pi} \Pi \theta + \alpha_0 \nabla_\mu n^\mu + \psi_{\alpha_{n\Pi}} n_\mu \dot{u}^\mu + \psi_{\alpha_{\Pi n}} n_\mu \nabla^\mu \alpha \right] \\ & - \frac{n^\mu}{T} \left[T \nabla_\mu \alpha - \beta_1 \dot{n}^\mu - \beta_{nn} n_\mu \theta + \alpha_0 \nabla_\mu \Pi + \alpha_1 \nabla_\nu \pi_\mu^\nu + \tilde{\psi}_{\alpha_{n\Pi}} \Pi \dot{u}_\mu \right. \\ & \left. + \tilde{\psi}_{\alpha_{\Pi n}} \Pi \nabla_\mu \alpha + \tilde{\chi}_{\alpha_{\pi n}} \pi_\mu^\nu \nabla_\nu \alpha + \tilde{\chi}_{\alpha_{n\pi}} \pi_\mu^\nu \dot{u}_\nu \right] \\ & + \frac{\pi^{\mu\nu}}{T} \left[\sigma_{\mu\nu} - \beta_2 \dot{\pi}_{\mu\nu} - \beta_{\pi\pi} \theta \pi_{\mu\nu} - \alpha_1 \nabla_{\langle\mu} n_{\nu\rangle} - \chi_{\alpha_{\pi n}} n_{\langle\mu} \nabla_{\nu\rangle} \alpha - \chi_{\alpha_{n\pi}} n_{\langle\mu} \dot{u}_{\nu\rangle} \right] \end{aligned} \quad (2.57)$$

where $\theta = \partial_\mu u^\mu$. The second law of thermodynamics $T \partial_\mu S^\mu \geq 0$ is guaranteed to be satisfied if we have,

$$T \partial_\mu S^\mu = \frac{\Pi^2}{\zeta} - \frac{n^\mu n_\mu}{\lambda} + \frac{\pi^{\mu\nu} \pi_{\mu\nu}}{2\eta}, \quad (2.58)$$

therefore, π , n^μ and $\pi^{\mu\nu}$ should satisfy the following equations,

$$\left[\theta + \beta_0 \dot{\Pi} + \beta_{\Pi\Pi} \Pi \theta + \alpha_0 \nabla_\mu n^\mu + \psi_{\alpha_{n\Pi}} n_\mu \dot{u}^\mu + \psi_{\alpha_{\Pi n}} n_\mu \nabla^\mu \alpha \right] = -\frac{\Pi}{\zeta}, \quad (2.59)$$

$$\begin{aligned} & \left[T \nabla_\mu \alpha - \beta_1 \dot{n}^\mu - \beta_{nn} n_\mu \theta + \alpha_0 \nabla_\mu \Pi + \alpha_1 \nabla_\nu \pi_\mu^\nu + \tilde{\psi}_{\alpha_{n\Pi}} \Pi \dot{u}_\mu \right. \\ & \left. + \tilde{\psi}_{\alpha_{\Pi n}} \Pi \nabla_\mu \alpha + \tilde{\chi}_{\alpha_{\pi n}} \pi_\mu^\nu \nabla_\nu \alpha + \tilde{\chi}_{\alpha_{n\pi}} \pi_\mu^\nu \dot{u}_\nu \right] = \frac{n^\mu}{\lambda}, \end{aligned} \quad (2.60)$$

$$\left[\sigma_{\mu\nu} - \beta_2 \dot{\pi}_{\mu\nu} - \beta_{\pi\pi} \theta \pi_{\mu\nu} - \alpha_1 \nabla_{\langle\mu} n_{\nu\rangle} - \chi_{\alpha_{\pi n}} n_{\langle\mu} \nabla_{\nu\rangle} \alpha - \chi_{\alpha_{n\pi}} n_{\langle\mu} \dot{u}_{\nu\rangle} \right] = \frac{\pi^{\mu\nu}}{2\eta}, \quad (2.61)$$

where $\lambda, \zeta, \eta \geq 0$ are the transport coefficients called coefficient of charge conductivity, bulk viscosity and shear viscosity respectively. Coefficients $\alpha_i, \beta_i, \alpha_{XY}, \beta_{XX}$ are the additional

transport coefficients and the parameters ψ, χ with $\tilde{\psi} = 1 - \psi$ and $\tilde{\chi} = 1 - \chi$ included the contributions due to the cross terms of Π and $\pi^{\mu\nu}$ with n^μ .

Onsager coefficients in this case too, can be obtained using the parameterization[see Eq.(2.25)],

$$\gamma_{11} = 2T \left[\eta \Delta^{\mu\alpha} \Delta^{\nu\beta} + \frac{1}{2} \left(\zeta - \frac{2}{3} \eta \right) \Delta^{\mu\nu} \Delta^{\alpha\beta} \right] \frac{1}{\Delta V}, \quad (2.62)$$

$$\gamma_{22} = -\frac{\lambda T \Delta^{\mu\nu}}{\Delta V}. \quad (2.63)$$

It should be noted that in Ref. [18] the authors have used $\frac{n^\mu}{\lambda}$ in Eq.(2.57) instead of $\frac{n^\mu}{\lambda T}$ and therefore Onsager coefficient differ by factor T (see for example, Eqs.(2.26 and 2.63)). The correlation functions can be written as,

$$\langle S_{vis}^{\mu\nu}(x_1) S_{vis}^{\alpha\beta}(x_2) \rangle = 2T \left[\eta (\Delta^{\mu\alpha} \Delta^{\nu\beta} + \Delta^{\mu\beta} \Delta^{\nu\alpha}) + \left(\zeta - \frac{2}{3} \eta \right) \Delta^{\mu\nu} \Delta^{\alpha\beta} \right] \delta(x_1 - x_2), \quad (2.64)$$

$$\langle I^\mu(x_1) I^\nu(x_2) \rangle = -2\lambda T \Delta^{\mu\nu} \delta(x_1 - x_2), \quad (2.65)$$

$$\langle S_{vis}^{\mu\nu}(x_1) I^\alpha(x_2) \rangle = 0. \quad (2.66)$$

2.3.3 DKR hydrodynamics

In Ref. [19], it was demonstrated that derivation of relativistic viscous hydrodynamic equation from the 14-moment method done by Israel and Stewart may not be unique. In Ref. [19], authors obtained relativistic dissipative hydrodynamic equations for the dissipative fluxes as,

$$\dot{\Pi} = -\frac{\Pi}{\tau_\Pi} - \beta_\Pi \theta - \delta_{\Pi\Pi} \Pi \theta + \lambda_{\Pi\pi} \pi^{\mu\nu} \sigma_{\mu\nu}, \quad (2.67)$$

$$\dot{\pi}^{\langle\mu\nu\rangle} = -\frac{\pi^{\mu\nu}}{\tau_\pi} + 2\beta_\pi \sigma^{\mu\nu} + 2\pi_\alpha^{\langle\mu} \omega^{\nu\rangle\alpha} - \delta_{\pi\pi} \pi^{\mu\nu} \theta - \tau_{\pi\pi} \pi_\alpha^{\langle\mu} \sigma^{\nu\rangle\alpha} + \lambda_{\pi\Pi} \Pi \sigma^{\mu\nu}. \quad (2.68)$$

It should be noted that Eq.(2.68) contains vorticity term ω . This arises due to kinetic theory considerations and such term does not appear in the MIS hydrodynamics [13]. Note that in writing the above equations we have considered the fluid with no net baryon number. Thus the Eq.(2.7) with no net baryon number can be written as,

$$\partial_\mu (s u^\mu) = \frac{\pi^{\mu\nu} \sigma_{\mu\nu}}{2T} - \frac{\Pi \Delta_\alpha u^\alpha}{T}, \quad (2.69)$$

$$\Delta_\alpha u^\alpha = -\frac{\dot{\Pi}}{\beta_\Pi} - \frac{\Pi}{\beta_\Pi \tau_\Pi} - \frac{\delta_{\Pi\Pi} \Pi \theta}{\beta_\Pi} + \frac{\lambda_{\Pi\pi} \pi^{\mu\nu} \sigma_{\mu\nu}}{\beta_\Pi}, \quad (2.70)$$

$$\sigma_{\mu\nu} = \frac{\dot{\pi}^{\langle\mu\nu\rangle}}{2\beta_\pi} + \frac{\pi^{\mu\nu}}{2\beta_\pi \tau_\pi} - \frac{\pi_\alpha^{\langle\mu} \omega^{\nu\rangle\alpha}}{\beta_\pi} + \frac{\delta_{\pi\pi} \pi^{\mu\nu} \theta}{2\beta_\pi} + \frac{\tau_{\pi\pi} \pi_\alpha^{\langle\mu} \sigma^{\nu\rangle\alpha}}{2\beta_\pi} - \frac{\lambda_{\pi\Pi} \Pi \sigma^{\mu\nu}}{2\beta_\pi}. \quad (2.71)$$

Assuming pressure gradient to be small and neglecting the higher (third) order term it is easy to show that $\dot{\pi}^{\langle\mu\nu\rangle} = \dot{\pi}^{\mu\nu}$.

Neglecting the bulk viscosity and using Eq.(2.71) in Eq.(2.69) one gets,

$$\begin{aligned}
\partial_\mu S^\mu = & \left[-\partial_\mu \left(\frac{u^\mu}{2\beta_\pi T} \right) + \frac{1}{2\beta_\pi \tau_\pi T} \right] \pi^{\alpha\beta} \pi_{\alpha\beta} \\
& - u^\beta \partial_\mu \left(\frac{\tau_{\pi\pi} \pi^{\alpha\mu} \pi_{\alpha\beta}}{4\beta_\pi T} \right) - u^\lambda \partial_\mu \left(\frac{\tau_{\pi\pi} \pi_\alpha^\mu \pi_\lambda^\alpha}{4\beta_\pi T} \right) + u^\mu \partial_\mu \left(\frac{\tau_{\pi\pi} \pi^{\alpha\beta} \pi_{\alpha\beta}}{6\beta_\pi T} \right) \\
& - u^\mu \partial_\mu \left(\frac{\delta_{\pi\pi} \pi^{\alpha\beta} \pi_{\alpha\beta}}{2\beta_\pi T} \right) + u^\beta \partial_\mu \left(\frac{\pi_{\alpha\beta} \pi^{\alpha\mu}}{2\beta_\pi T} \right) - u^\lambda \partial_\mu \left(\frac{\pi_\alpha^\mu \pi_\lambda^\alpha}{2\beta_\pi T} \right), \tag{2.72}
\end{aligned}$$

where, S^μ is the non-equilibrium entropy current for DKR hydrodynamics and has the form,

$$\begin{aligned}
S^\mu = & \left(su^\mu - \frac{\pi^{\alpha\beta} \pi_{\alpha\beta} u^\mu}{4\beta_\pi T} - \frac{\delta_{\pi\pi} \pi^{\alpha\beta} \pi_{\alpha\beta} u^\mu}{2\beta_\pi T} - \frac{\tau_{\pi\pi} \pi^{\alpha\mu} \pi_{\alpha\beta} u^\beta}{4\beta_\pi T} - \frac{\tau_{\pi\pi} \pi_\alpha^\mu \pi_\lambda^\alpha u^\lambda}{4\beta_\pi T} \right. \\
& \left. + \frac{\tau_{\pi\pi} \pi^{\alpha\beta} \pi_{\alpha\beta} u^\mu}{6\beta_\pi T} + \frac{\pi_{\alpha\beta} \pi^{\alpha\mu} u^\beta}{2\beta_\pi T} - \frac{\pi_\alpha^\mu \pi_\lambda^\alpha u^\lambda}{2\beta_\pi T} \right). \tag{2.73}
\end{aligned}$$

Since $\pi^{\mu\nu} \ll T^{\mu\nu}$, so $O((\pi^{\mu\nu})^2)$ terms will be smaller and its divergence will be much smaller. So on right hand side of Eq.(2.72) one can drop the divergence terms. Similarly, one can neglect the terms having divergence of four-velocity multiplied by $O((\pi^{\mu\nu})^2)$. Thus we can write Eq.(2.72) as,

$$\frac{dS}{dt} = \int d^3x \frac{\pi^{\alpha\beta}}{T} \left[\left(\frac{1}{2\beta_\pi \tau_\pi} \right) \right] \pi_{\alpha\beta}. \tag{2.74}$$

A comparison of the above expression with the phenomenological equation (Eq.(2.1)) yields,

$$\dot{x} \rightarrow \pi^{\alpha\beta}, \quad X \rightarrow -\frac{1}{T} \left[\left(\frac{1}{2\beta_\pi \tau_\pi} \right) \right] \pi_{\alpha\beta} \Delta V. \tag{2.75}$$

Again by comparing Eq.(2.75) with Eq.(2.2)(when $\xi = 0$) one can get,

$$\gamma X = -\pi^{\mu\nu}. \tag{2.76}$$

By considering γ to be of the form,

$$\gamma = \left[A \Delta^{\mu\nu\alpha\beta} + B \Delta^{\mu\nu} \Delta^{\alpha\beta} \right] \frac{1}{\Delta V}, \tag{2.77}$$

and using Eq.(2.77) one finds that $A = \frac{T}{\left(\frac{1}{2\beta_\pi \tau_\pi} \right)}$ and $B = 0$. Thus the viscous correlations are,

$$\langle S_{vis}^{\mu\nu}(x_1) S_{vis}^{\alpha\beta}(x_2) \rangle = 2T\beta_\pi \tau_\pi \left(\Delta^{\mu\alpha} \Delta^{\nu\beta} + \Delta^{\mu\beta} \Delta^{\nu\alpha} - \frac{2}{3} \Delta^{\mu\nu} \Delta^{\alpha\beta} \right) \delta(x_1 - x_2). \tag{2.78}$$

2.3.4 Conformal viscous hydrodynamics

The entropy current for the conformal hydrodynamics [20] can be written as,

$$S^\mu = \left(su^\mu - \frac{\tau_\Pi}{4\eta T} \Pi_{\alpha\beta} \Pi^{\alpha\beta} u^\mu \right). \quad (2.79)$$

One can easily find the following expression for the Onsager coefficient and the correlation function,

$$\gamma = 2\eta T \left[\Delta^{\mu\alpha} \Delta^{\nu\beta} - \frac{1}{3} \Delta^{\mu\nu} \Delta^{\alpha\beta} \right] \frac{1}{\Delta V}, \quad (2.80)$$

$$\langle S_{vis}^{\mu\nu}(x_1) S_{vis}^{\alpha\beta}(x_2) \rangle = 2\eta T \left[(\Delta^{\mu\alpha} \Delta^{\nu\beta} + \Delta^{\mu\beta} \Delta^{\nu\alpha}) - \frac{2}{3} \Delta^{\mu\nu} \Delta^{\alpha\beta} \right] \delta(x_1 - x_2). \quad (2.81)$$

3. Calculation of correlation functions in boost-invariant Hydrodynamics

As an example we apply the results obtained in the previous sections to the relativistic heavy-ion collisions for the Bjorken scaling solution.

According to Bjorken scenario in heavy ion collisions, the reaction volume is strongly expanded in the longitudinal direction, i.e along the collision axis(z-axis). So one can assume that there is no transverse flow. Thus one can describe flow in 1 + 0 dimension [22]. It is useful to introduce the light cone variable y and proper time τ which are defined by,

$$\tau = \sqrt{t^2 - z^2} \text{ and } y = \text{arc tanh}(z/t) = \frac{1}{2} \ln\left(\frac{t+z}{t-z}\right). \quad (3.1)$$

The partial derivatives in time and space can be expressed as,

$$\begin{bmatrix} \partial_t \\ \partial_z \end{bmatrix} = \begin{bmatrix} \cosh y & -\sinh y \\ -\sinh y & \cosh y \end{bmatrix} = \begin{bmatrix} \partial_\tau \\ \frac{1}{y} \partial_y \end{bmatrix}. \quad (3.2)$$

The flow velocity under the scaling assumption can be written as, $u^\mu = \gamma(1, 0, 0, v_z) = (\frac{t}{\tau}, 0, 0, \frac{z}{\tau}) = (\cosh y, 0, 0, \sinh y)$. We consider only longitudinal flow fluctuations and parameterize the flow velocity [23] as,

$$u^\mu = (\cosh \bar{\theta}, \sinh \bar{\theta}), \quad (3.3)$$

where $\bar{\theta} = y + \delta\bar{\theta}(y, \tau)$ and $\delta\bar{\theta}(y, \tau)$ are the fluctuations in the longitudinal flow. In Bjorken scaling limit, $\bar{\theta} = y$. With this parameterization and using the transformation of derivatives one can introduce the operators D, ∇ such that,

$$\begin{bmatrix} D \\ \nabla \end{bmatrix} = \begin{bmatrix} \cosh(\bar{\theta} - y) & \sinh(\bar{\theta} - y) \\ \sinh(\bar{\theta} - y) & \cosh(\bar{\theta} - y) \end{bmatrix} = \begin{bmatrix} \partial_\tau \\ \frac{1}{y} \partial_y \end{bmatrix}. \quad (3.4)$$

In the scaling limit, $D = u^\mu \partial_\mu = \frac{\partial}{\partial \tau} = \partial_\tau$ and $\partial_\mu u^\mu = \nabla \bar{\theta} = \frac{1}{\tau}$.

Since $S^{\mu\nu}$ satisfies the condition,

$$u_\mu S^{\mu\nu} = 0. \quad (3.5)$$

One can write $S^{\mu\nu}$ as [6],

$$S^{\mu\nu} = w(\tau) f(y, \tau) \Delta^{\mu\nu}, \quad (3.6)$$

where, $w = \epsilon + p = Ts$ and f is a dimensionless quantity which satisfy $\langle f \rangle = 0$, where $\langle \rangle$ denotes the ‘average value’. In heavy-ion collision experiments at LHC or RHIC a baryon free quark-gluon plasma is expected to be produced, therefore $q^\mu = 0$. Thus only viscous-correlations are of interest, which for MIS, JBP, third order and conformal case can be written as,

$$\langle f(y_1, \tau_1) f(y_2, \tau_2) \rangle = \frac{2T(\tau_1)}{A\tau_1 w^2(\tau_1)} \left[\frac{4}{3} \eta(\tau_1) + \zeta(\tau_1) \right] \delta(\tau_1 - \tau_2) \delta(y_1 - y_2), \quad (3.7)$$

where $\delta(x_1 - x_2)_{Transverse}$ is replaced by effective transverse area A of colliding nuclei. Note that these correlations are same as that obtained by authors in Ref. [6] for Navier-Stokes case. Similarly, for DKR case one can write the viscous correlations as,

$$\langle f(y_1, \tau_1) f(y_2, \tau_2) \rangle = \frac{2T(\tau_1) \left(\frac{4}{3} \beta_\pi \tau_\pi \right)}{A\tau_1 w^2(\tau_1)} \delta(\tau_1 - \tau_2) \delta(y_1 - y_2). \quad (3.8)$$

Defining $\eta = \beta_\pi \tau_\pi$ (see Ref. [19]) and neglecting the bulk viscosity for the correlation functions, one can rewrite the correlations for all the models of hydrodynamics as,

$$\langle f(y_1, \tau_1) f(y_2, \tau_2) \rangle = \frac{X(\tau_1)_{[E]}}{A} \delta(\tau_1 - \tau_2) \delta(y_1 - y_2), \quad (3.9)$$

where,

$$X(\tau_1)_{[E]} = \frac{8}{3\tau_1 w(\tau_1)} \left(\frac{\eta(\tau_1)}{s(\tau_1)} \right)_{[E]}. \quad (3.10)$$

Here, subscript $[E]$ denotes the particular type of hydrodynamics model considered from the set of hydrodynamics models, for example $[E] = [MIS, JBP, DKR, NS, \dots]$.

It is useful to study the correlation function normalized by the initial value of the correlation obtained using the Navier-Stokes theory i.e. $C(\tau) = \frac{w^2(\tau) X(\tau)_{[E]}}{w^2(\tau_0)_{NS} X(\tau_0)_{NS}}$ where, τ_0 is the initial-time for the hydrodynamics. $C(\tau)$ can be written as,

$$C(\tau)_{[E]} = \frac{\left(\frac{\tau_0}{\tau_1} \right) \left(\frac{\eta(\tau)}{s(\tau)} \right)_{[E]} \frac{w(\tau)}{w(\tau_0)}}{\left(\frac{\eta(\tau)}{s(\tau)} \right)_{NS}}. \quad (3.11)$$

Further, we neglect the effect of bulk-viscosity by considering the initial temperature T_i to be much larger than the critical temperature, $T_c = 0.190$ GeV. Now, in the Landau-Lifshitz frame, the energy and the momentum conservation laws are given by,

$$u_\nu \partial_\mu T^{\mu\nu} = D\epsilon + (\epsilon + p) \nabla \bar{\theta} - \pi_{\mu\nu} \nabla^{(\mu} u^{\nu)} - S_{\mu\nu} \nabla^{(\mu} u^{\nu)} = 0, \quad (3.12)$$

$$\Delta_\nu^\alpha \partial_\mu T^{\mu\nu} = (\epsilon + p) D u^\alpha - \nabla^\alpha p + \Delta^{\alpha\nu} \nabla^\sigma \pi_{\nu\sigma} - \pi^{\alpha\nu} D u_\nu + \Delta^{\alpha\nu} \partial^\sigma S_{\sigma\nu} = 0, \quad (3.13)$$

where, $\pi^{\alpha\beta}$ is the shear stress tensor and will be different for different models of hydrodynamics (e.g, MIS, JBP, DKR, NS etc.).

In the scaling limit $\bar{\theta} = y$, $D = u^\mu \partial_\mu = \partial_\tau$, $\partial_\mu u^\mu = \nabla \bar{\theta} = \frac{1}{\tau}$ and shear stress tensor $\pi^{\mu\nu}$ has only diagonal components i.e $\pi^{\mu\nu} = \text{diag}(0, \pi/2, \pi/2, -\pi)$.

Therefore, one can write the above equations as [19],

$$\partial_\tau \epsilon = -\frac{(\epsilon + p)}{\tau} + \frac{\pi}{\tau}. \quad (3.14)$$

Here, the noise term is considered to be smaller than the background quantities.

Equation for π in the scaling limit, for MIS, DKR and JBP hydrodynamics can be written as,

$$\partial_\tau \pi + \frac{\pi}{\tau} = \beta_\pi \frac{4}{3\tau} - \lambda \frac{\pi}{\tau}. \quad (3.15)$$

First we consider MIS case. In this literature on relativistic heavy-ion collisions $\lambda = 0$ case is being considered [15, 21]. The coefficients β_π and τ_π can be written as,

$$\beta_\pi = \frac{2p}{3}, \quad \tau_\pi^{-1} = \frac{5}{9} \frac{\sigma p}{T}. \quad (3.16)$$

For JBP models parameters β_π , τ_π and λ are as follows,

$$\beta_\pi = \frac{2p}{3}, \quad \tau_\pi^{-1} = \frac{5}{9} \frac{\sigma p}{T}, \quad \lambda = 4/3, \quad (3.17)$$

where, σ is the total cross-section [19] and it is assumed to be independent of energy [17, 24]. In the massless gas limit the expressions for the coefficients β_π and τ_π are same in both JBP and MIS dynamics [18]. For DKR hydrodynamics, the parameters β_π , τ_π [19] and λ [25] are,

$$\beta_\pi = \frac{4p}{5}, \quad \tau_\pi^{-1} = \frac{3}{5} \frac{\sigma p}{T}, \quad \lambda \equiv \frac{1}{3} \tau_{\pi\pi} + \delta_{\pi\pi} = \frac{38}{21}. \quad (3.18)$$

In what follows we consider the ideal equation of state, $\epsilon = 3p$ and the pressure is given by the bag model, $p = \frac{\pi^2}{30} T^4$. Further, we consider the initial values for temperature $T_i = 0.310$ GeV and viscous stress $\pi = 0$ for the causal hydrodynamics and numerically solve Eqs. (3.14, 3.15) for evaluating the correlations (3.11) in case of MIS, JBP, DKR hydrodynamics. However, for the Navier-Stokes hydrodynamics one needs to solve only Eq. (3.14) with same value of initial temperature and the viscous stress is given by,

$$\pi = \eta \frac{4}{3\tau}. \quad (3.19)$$

The results of the numerical work are presented in the following section.

4. Results and Discussions

We have studied fluctuations in various models of relativistic causal viscous hydrodynamics. Eqs. (2.27-2.29), (2.47-2.49), (2.56), (2.64-2.66), (2.78) and (2.81) represent our main results describing the correlation functions for various models of relativistic causal hydrodynamics. First we should like to note here that the form of the correlation functions given by Eqs. (2.27-2.29), (2.47-2.49), (2.56), (2.64-2.66), (2.78) and (2.81) are strikingly similar to the correlation functions obtained using relativistic Navier-Stokes theory [3, 6]. It should be noted that Navier-Stokes equations and the other relativistic causal hydrodynamics theories are consistent with the condition that four-divergence of the entropy-current $T \partial_\mu S^\mu$ remains




Figure 1: (a), (b) and (c) show time evolution of the correlation functions for NS, MIS, DKR and JBP hydrodynamics for the initial temperature $T_i = 310 \text{ MeV}$. The coefficient of viscosity is calculated using $\eta_{DKR} = \frac{4T}{3\sigma}$. The scaling $\eta_{MIS} = \eta_{JBP} = 9/10\eta_{DKR}$ and $\eta_{NS} = \frac{7.59}{8}\eta_{DKR}$ ensure that the cross-section remains same in the comparison between the models of hydrodynamics. Case (a) corresponds to $\frac{\eta_{DKR}}{s} = 0.08$, while cases (b), (c) depicts for $\frac{\eta_{DKR}}{s} = 0.24$, $\frac{\eta_{DKR}}{s} = 1.60$ respectively. In all the figures, solid (red), dotdashed (blue), dotted (green) and dashed (purple) curves correspond to the NS, MIS, DKR and JBP Hydrodynamics respectively.

always positive. The expression for divergence of the entropy four-currents (see Eq. 2.17) contains tensors like $\pi_{\mu\nu}$, q^μ and scalar Π . Since the derivation of the Onsager coefficients γ uses $\Delta_{\mu\nu}\pi^{\mu\nu} = 0$, $q_\mu u^\mu = 0$ and $u_\mu\pi^{\mu\nu} = 0$ as constraints on the dissipative fluxes, the form of the correlation function (2.27-2.29), (2.47-2.49), (2.56), (2.64-2.66), (2.78) and (2.81) remains similar to the one obtained using the relativistic Navier-Stokes theory. But in case of DTT kind of hydrodynamics, it is not clear if divergence of the entropy four-current can be expressed directly in terms of scalar product of the viscosity and heat-flux tensors.

Since forms of the equations of relativistic causal hydrodynamics of MIS, DKR, JBP are different, the space-time evolution of the correlation functions can be very different from the evolution of the correlations in the Navier-Stokes theory. Next, we have shown that when one considers the third order correction [17] to the MIS hydrodynamics, the form of correlation function remains to be same as the one obtained using the MIS theory.




Figure 2: (a), (b) and (c) show the time evolution of the correlation functions for NS, MIS, DKR and JBP hydrodynamics. For Fig.(2a), the ratio of the viscosity to entropy density $\frac{\eta}{s} = 0.08$ is kept same for all the four cases, while figure (2b) and (2c) correspond to cases with $\frac{\eta}{s} = 0.24$ and $\frac{\eta}{s} = 1.60$ respectively. Here, solid (red), dotdashed (blue), dotted (green) and dashed (purple) curves correspond to the NS, MIS, DKR and JBP Hydrodynamics respectively for all the figures.

Further, we have considered the case of conformal hydrodynamics when the bulk-viscosity is absent. In this case also the form of the Onsager coefficients and the viscous correlations remains same as that of MIS theory. Again the temporal evolution of the correlations may be different than the MIS theory as the underlying hydrodynamical equations are different.

As there exists a variety of the relativistic causal hydrodynamics models, it is natural to ask the question that which kind of hydrodynamics to be applied for a given physical situation. This question is rather hard to answer in general. In order to understand the evolution of the correlation functions in some details we have calculated the normalized correlation functions given by Eqn.(3.11) for an expanding one-dimensional boost-invariant (Bjorken) flow. In figures (1-2), we plot the normalized correlation function $C(\tau)$ (Eqn.3.11) for MIS, JBP, DKR and NS hydrodynamics as a function of time τ . The initial values of the temperature, $T_i = 310 \text{ MeV}$ and the viscous stress $\pi = 0$ are kept same for all the cases shown in figures (1-2). Each figure has curves representing the four different models

of hydrodynamics: the solid line describes the correlation function from the Navier-Stokes equations, the dotted-dashed curve represents the case of MIS equations, the dotted line describes the DKR case, while the dashed curve describes JBP case. There are two kind of comparisons possible between the correlation functions for the different models of relativistic viscous hydrodynamics. In one such comparison the energy-independent cross-section[see Eqns. (3.16-3.18)] is kept same for all the different versions of the hydrodynamics [19]. Following [19,24] one can write the relation between the viscosity coefficient for the different models of hydrodynamics as $\eta_{DKR} = \frac{4T}{3\sigma}$, $\eta_{MIS} = \frac{6T}{5\sigma} = \eta_{JBP}$ and $\eta_{NS} = 0.8436 \frac{3T}{2\sigma}$. The scaling $\eta_{MIS} = \eta_{JBP} = 9/10 \eta_{DKR}$ and $\eta_{NS} = \frac{7.59}{8} \eta_{DKR}$ ensure that the cross-section remains same in the comparison between the different models of hydrodynamics. In another way of comparing the correlation-functions, the ratio of viscosity coefficient to the entropy density η/s is kept same for the different models of the hydrodynamics, while the transport cross-section is varied for the different hydrodynamics models.

Figure (1) corresponds to the case when the cross-section is kept same for all the models of the hydrodynamics. The case(a) corresponds to $\frac{\eta_{DKR}}{s} = 0.08$, while η/s for the other hydrodynamics models is calculated using the scaling. In this case, all the four curves overlap with each other. This case corresponds to the *nearly ideal* hydrodynamics. The case (b) and (c) corresponds to $\frac{\eta_{DKR}}{s} = 0.24$ and $\frac{\eta_{DKR}}{s} = 1.6$ respectively. One can notice here that the correlation function corresponding to the Navier-Stokes hydrodynamics consistently gives higher values of the correlation for values of $\frac{\eta_{DKR}}{s} > 0.08$. The DKR curve also gives higher value of the correlation in comparison with JBP and MIS hydrodynamics. The figure inset in case(b) shows the plots of DKR, JBP and MIS correlation functions in τ range between $5fm/c$ to $7fm/c$. The reason why the MIS and JBP dynamics follows each other is simple as values of parameters β_π and τ_π are same for both the cases and dynamics of viscous stress is governed by the Eq.(3.15). In case (c) one can see that there is a significant difference between the Navier-Stokes equation and the other causal hydrodynamics models. The Navier-Stokes correlation rises for such a high value of the viscosity. This is an unphysical behavior. It is well-known that when the Navier-Stokes equation is applied to one-dimensional boost-invariant flow, temperature of the flow can increase for some time and reach a maximum value before it starts to plummet [26].

Figure (2) corresponds to the case when the ratio of the viscosity coefficient to the entropy density is kept same for all the four models of hydrodynamics. Case (2a) corresponds to the situation when $\frac{\eta}{s} = 0.08$ for all four models of the hydrodynamics. In this case, when the viscosity is very low, all the correlations $C(\tau)$ for all the four models (NS, MIS, DKR and JBP) of hydrodynamics overlap. Figure (2b) corresponds to the case when $\frac{\eta}{s} = 0.24$. Here, again the Navier-Stokes hydrodynamics consistently gives higher values of the correlation function. The inset diagram in figure (2b) shows the plots of DKR, JBP and MIS correlation functions in τ range between $5fm/c$ to $7fm/c$. Figure (2c) corresponds to case when $\frac{\eta}{s} = 1.6$, here the Navier-Stokes dynamics gives the unphysical behavior.

The cases studied in figures (1-2) show that Navier-Stokes hydrodynamics gives consistently higher values for the correlation. The difference between the correlation functions obtained using the Navier-Stokes dynamics and of the correlation functions from the other causal approaches (MIS, DKR and JBP) increases with increasing values of η/s . However,

the differences between the correlation-functions of different causal hydrodynamics are not very significant. This could be due to the fact that the present example deals with the boost-invariant one-dimensional flow. In a non-central heavy-ion collision, the vorticity in the flow can play a significant role [27]. As one can notice from Eq.(2.68) that vorticity can drive dynamics of the viscous stress in DKR hydrodynamics. However this will require to solve hydrodynamical equation in $2 + 1$ or $3 + 1$ dimensions. This is at present, beyond the scope of this work.

Thus we have shown using one-dimensional flow as an example that evolution of the correlation functions of fluctuations can sensitively depend on the model of hydrodynamics. The well-separated nature of the correlation-function of the relativistic Navier-Stokes equations in comparison with the models of the causal hydrodynamics may allow us, we believe, in figuring out which version of the relativistic dissipative hydrodynamics might be suitable for describing the heavy-ion collision dynamics.

5. Conclusions

We have studied fluctuations in various models of relativistic causal hydrodynamics. We have found that the general properties of the dissipative part of the energy-momentum tensor due to the viscosity and heat-flux play an important role in determining the Onsager coefficients and the correlation functions. We find that all the models of causal relativistic hydrodynamics have similar form of the correlation function. In addition, we have applied these results to one-dimensional boost-invariant hydrodynamic flow to investigate the temporal evolution of the correlation functions for the different hydrodynamics models. We believe that our study may help in choosing a proper version of the causal hydrodynamics for a given physical situation.

Note added: After this manuscript was prepared, we have found in Ref [28] on arXiv that the authors have applied the fluctuation-dissipation relation to the relativistic viscous hydrodynamics with the memory effects.

6. Acknowledgements

The authors (AK, JRB) would like to thank to Prof. P. K. Kaw.

References

- [1] L. D. Landau and E. M. Lifshitz, *Statistical Physics: Part 1* (Pergamon, Oxford, 1980).
- [2] L. D. Landau and E. M. Lifshitz, *Statistical Physics: Part 2* (Pergamon, Oxford, 1980).
- [3] N. Salie, R. Wuffert, and W. Zimdahl, *Fluctuations in the general relativistic theory of fluids and solids*, J. Phys. A: Math. Gen. **16**, 3533 (1983).
- [4] E. Calzetta, *Relativistic fluctuating hydrodynamics*, Class. Quantum Grav. **15**, 653 (1998).
- [5] R. Geroch, and L. Lindblom, *Dissipative relativistic fluid theories of divergence type*, Phys. Rev. **D41**, 1855 (1990).

- [6] J. I. Kapusta, B. Muller, and M. Stephanov, *Relativistic theory of hydrodynamic fluctuations with applications to heavy-ion collisions*, Phys. Rev. **C85**, 054906 (2012).
- [7] J. I. Kapusta, J. M. Torres-Rincon, *Thermal Conductivity and Chiral Critical Point in Heavy Ion Collisions*, Phys. Rev. **C86**, 054911 (2012).
- [8] W. A. Hiscock, and L. Lindblom, *Generic instabilities in first-order dissipative relativistic fluid theories*, Phys. Rev. **D31**, 725 (1985), and *Stability and causality in dissipative relativistic fluids*, Ann. Phys. **151**, 466 (1983).
- [9] P. Romatschke, *New developments in Relativistic Viscous Hydrodynamics*, Int. J. Mod. Phys. **E19**, 1 (2010) (arXiv:0902.3663[hep-ph]).
- [10] I. Müller, *To the Paradox of Heat Conduction Theory*, Z. Phys. **198**, 329 (1967); W. Israel, *Non-stationary Irreversible Thermodynamics: A Causal Relativistic Theory*, Ann. Phys. **100**, 310 (1976); W. Israel, and J. M. Stewart, *Transient Relativistic Thermodynamics and Kinetic Theory*, Ann. Phys. **118**, 341 (1979).
- [11] D. Jou, J. Casas-Vazquez, and G. Lebon, *Extended irreversible thermodynamics*, Rep. Prog. Phys. **51**, 1105 (1988); *ibid*, *Extended irreversible thermodynamics revisited (1988-98)*, **62**, 1035 (1999).
- [12] L. Herrera and D. Pavon, *Why hyperbolic theories of dissipation cannot be ignored: Comments on a paper by Kostadt and Liu*, Phys. Rev. **D64**, 088503 (2001); L. Herrera and D. Pavon, *Hyperbolic theories of dissipation: Why and when do we need them?*, Physica **A307**, 121 (2002).
- [13] A. Muronga, *Causal theories of dissipative relativistic fluid dynamics for nuclear collisions*, Phys. Rev. **C69**, 034903 (2004).
- [14] A. Muronga and D. H. Rischke, *Evolution of Hot, Dissipative Quark Matter in Relativistic Nuclear Collisions*, arXiv:nucl-th/0407114.
- [15] U. Heinz, H. Song, and A. K. Chaudhuri, *Dissipative hydrodynamics for viscous relativistic fluids*, Phys. Rev. **C73**, 034904 (2006); U. Heinz, *Viscous relativistic hydrodynamics*, arXiv:nucl-th/0512049.
- [16] M. Bastero-Gil, A. Berera, R. Cerezo, R. O. Ramos, and G. S. Vicente, *Stability analysis for the background equations for inflation with dissipation and in a viscous radiation bath*, Jour. Cos. Astro. Phys. **1211**, 042 (2012).
- [17] A. El, Z. Xu, and C. Greiner, *Extension of relativistic dissipative hydrodynamics to third order*, Phys. Rev. **C81**, 041901(R) (2010).
- [18] A. Jaiswal, R. S. Bhalerao, and S. Pal, *Complete relativistic second-order dissipative hydrodynamics from the entropy principle*, Phys. Rev. **C87**, 021901(R) (2013).
- [19] G. S. Denicol, T. Koide, and D. H. Rischke, *Dissipative relativistic fluid dynamics: a new way to derive the equations of motion from kinetic theory*, Phys. Rev. Lett. **105**, 162501 (2010).
- [20] R. Baier, P. Romatschke, D. T. Son et al., *Relativistic viscous hydrodynamics, conformal invariance, and holography*, J. High Energy Phys. **0804** (2008) 100.
- [21] A. Muronga, *Relativistic dynamics of nonideal fluids: Viscous and heat-conducting fluids. I. General aspects and 3+1 formulation for nuclear collisions*, Phys. Rev. **C76**, 014909 (2007).
- [22] J. D. Bjorken, *Highly relativistic nucleus-nucleus collisions: The central rapidity region*, Phys. Rev. **D27**, 140 (1983).

- [23] H. Kouno, M. Maruyama, and F. Takagi, *Relativistic Hydrodynamics of Quark-Gluon Plasma and Stability of Scaling Solutions*, Phys. Rev. **D41**, 2903 (1990).
- [24] P. Huovinen and D. Molnar, *The applicability of causal dissipative hydrodynamics to relativistic heavy-ion collision*, Phys. Rev. **C79**, 014906 (2009), and *Applicability of viscous hydrodynamics at RHIC*, Nucl. Phys. **A830**, 475C (2009).
- [25] A. Jaiswal, Chapman-Enskog expansion and relativistic dissipative hydrodynamics, arXiv:nucl-th/1302.6311.
- [26] R. Baier, P. Romatschke and U. A. Wiedemann, *Dissipative hydrodynamics and heavy-ion collisions*, Phys. Rev. **C73**, 064903 (2003).
- [27] S. Florschinger, and U. A. Wiedemann, *Fluctuations around Bjorken Flow and the onset of turbulent phenomena*, J. High Energy Phys. **11** (2011) 100.
- [28] K. Murase and T. Hirano, *Relativistic fluctuating hydrodynamics with memory functions and colored noises*, arXiv:nucl-th/1304.3243.